



Core Imaging Facility

Ben Fowler, Manager
(405) 271-7545
<http://Imaginglinux2.omrf.ouhsc.edu/>

Oklahoma Medical Research Foundation

825 N.E. 13th Street
Oklahoma City, OK 73104
405-271-6673
800-522-0211
<http://OMRF.org/>

Brief Immunogold Protocol **(Non-conjugated primary and conjugated secondary)**

1. Prepare a parafilm template using a long strip of parafilm. ****Do not touch the surface of the parafilm with your bare hands prior to staining.****
2. Lay down drops of PBS and hydrate grids until ready for next step (~5min).
4. Block with 2% BSA (or blocking solution of choice) for 30 minutes. ****May also use 3-5% NFDM or 2° Ab animal serum.****
5. 1 Ab (or PBS for control) for 30-45 minutes. Use at least a 30µl droplet, you may also cover grids and humidify.
6. Rinse 4X, 5 min. each with PBS or dH₂O.
7. 2 Ab** (or PBS for control) for 45 min -1 hour. (α gold; α protein A or G).
8. Rinse 3X, 5 min. each with dH₂O.
9. Post-fix if necessary with 2% PFA 10-15 min.
10. Air dry.
11. Store in grid box.

* Each set of experiments should contain a negative control (the slide is subjected to the same procedure, but instead of adding antibodies all steps receive buffer) and a 2 Ab control (no primary is added, but 2 Ab is added.) If desired, a non-specific primary antibody control may also be performed.

Controls you must have are PBS only; 2° only; and 1°, 2°. You may also have non-1°, 2°; 1°, non-2°, and positive control.